

Magnetospheric particle acceleration and X-ray emission of pulsars

Sevinç O. Tagieva¹ *, Aşkın Ankay² †, Arzu Mert Ankay² ‡

¹Academy of Science, Physics Institute,
Baku 370143, Azerbaijan Republic

²Boğaziçi University, Physics Department,
34342-Bebek, İstanbul, Turkey

Abstract

The available data on isolated X-ray pulsars, their wind nebulae, and the supernova remnants which are connected to some of these sources are analyzed. It is shown that electric fields of neutron stars tear off charged particles from the surface of neutron star and trigger the acceleration of particles. The charged particles are accelerated mainly in the field of magneto-dipole radiation wave. Power and energy spectra of the charged particles depend on the strength of the magneto-dipole radiation. Therefore, the X-ray radiation is strongly dependent on the rate of rotational energy loss and weakly dependent on the electric field intensity. Coulomb interaction between the charged particles is the main factor for the energy loss and the X-ray spectra of the charged particles.

Key words: Pulsar, PWN, X-ray

*e-mail:physic@lan.ab.az

†e-mail:askin.ankay@boun.edu.tr

‡arzu.mert@boun.edu.tr

1 Introduction

The number of isolated X-ray pulsars and the radio pulsars which have been detected in X-ray with characteristic age (τ) up to 10^6 yr located at distances less than 7 kpc from the Sun together with 2 pulsars in Magellanic Clouds is 35 (see Table 1). The so called magnetars (anomalous X-ray pulsars and soft gamma repeaters) and dim radio quiet neutron stars like the isolated young X-ray pulsar 1E 1207.4-5209 are not included in this list as our main aim in this article is to examine the non-thermal synchrotron (power-law) emission of isolated young pulsars with relatively well known ages based on the existing observational data. Also, our list includes only the pulsars which are certainly not accreting sources. In order to exclude thermal emission of pulsars, we will particularly focus our attention on the hard X-ray emission of the pulsars in the 2-10 keV band. On the other hand, we have included the old recycled millisecond pulsars in the list and examined them in detail throughout the text to make comparisons between them and the isolated young pulsars.

Most of the pulsars displayed in Table 1 have been observed in 0.1-2.4 keV band ^{1,2} and in 2-10 keV band ³ and all the available data have been collected, revised and put together as represented in the table. All of these pulsars have rate of rotational energy loss $\dot{E} > 3 \times 10^{34}$ erg/s except J1718-3718 for which $\dot{E}$ is as low as 1.6×10^{33} erg/s. Nine of these 35 pulsars are connected to F-type supernova remnants (SNRs) and 9 of them are connected to C-type SNRs. Therefore, there exist pulsar wind nebulae (PWNe) near these 18 pulsars which are located up to 7 kpc from the Sun. Four or five of the pulsars are connected to S-type SNRs (J0538+2817 – G180.0-1.7 connection is uncertain) and 2 other pulsars are connected to SNRs for which the morphological type is not known. Ten pulsars with detected X-ray emission are not connected to SNRs and 6 of them have characteristic ages $2 \times 10^5 < \tau < 10^6$ yr. Among the 35 pulsars, 2 of them are X-ray pulsars from which radio emission has not been observed and they are connected to SNRs (pulsar J1846-0258 to SNR G29.7-0.3 and pulsar J1811-1925 to SNR G11.2-0.3). Radio pulsar J1646-4346 ($d=4.51$ kpc) which is connected to C-type SNR G341.2+0.9 but not observed in X-ray is also included in the list. Other than these sources, we have also included 14 radio pulsars with $\dot{E} > 4 \times 10^{35}$ erg/s, $\dot{P} > 10^{-15}$ s/s and $d \leq 7$ kpc in Table 1. Also, 6 isolated millisecond pulsars from which X-ray radiation have been observed are displayed in Table 1.

Our main aims in this work are to analyze the conditions which are necessary for the acceleration of relativistic particles, the X-ray radiation produced in the magnetospheres of pulsars, and the presence or absence of PWNe around these objects.

2 Dependence of the ejection of relativistic particles on different parameters of pulsars

As known, very young pulsars have the capability to eject relativistic particles. Power and spectra of the particles must be related to different parameters of pulsars, particularly to the induced electric field intensity for which it is practically not possible to find an exact expression as there are significant uncertainties in the magnetic field structures of neutron stars in plasma. Therefore, we use the simplified relation

$$E_{el} = \frac{B_p R}{P} \quad (1)$$

⁴ which must be correct in the case when the braking of pulsar's rotation is connected to magneto-dipole radiation only. Here, R is the radius, B_p is the magnetic field intensity at the magnetic pole, and P is the rotation period of the neutron star. The number and energy spectra of the relativistic particles also depend on the conditions in the atmospheres of neutron stars (density and temperature). Independent of the mechanism, the rate of rotational energy loss ($\dot{E}$) depends only on the moment of inertia (I), the rotation period P and the time derivative of P ($\dot{P}$)

$$\dot{E} = \frac{4\pi^2 I \dot{P}}{P^3}. \quad (2)$$

If we assume that practically the rate of rotational energy loss is related only to the magneto-dipole radiation ^{4,5}, then the effective magnetic field $B = B_p \sin \alpha$ (where α is the angle between the surface dipole magnetic field axis and the rotation axis) and

$$\dot{E} = \frac{2}{3} \frac{\mu_{\perp}^2 \omega^4}{c^3} = \frac{8\pi^4}{3c^3} \frac{B^2 R^6}{P^4} \quad (3)$$

where $\mu_{\perp} = BR^3/2$ is the component of the magnetic moment perpendicular to the rotation axis, $\omega = 2\pi/P$ the angular velocity and c the speed of light. From expressions (1) and (3) we get

$$\dot{E} = \frac{8\pi^4}{3c^3} \frac{R^4}{P^2} E_{el}^2 \sin^2 \alpha. \quad (4)$$

As seen from this equation, $\dot{E}$ is directly proportional to E_{el}^2 and inversely proportional to P^2 .

From expressions (1)-(3), we can get

$$E_{el} \sim I^{1/2} R^{-2} \tau^{-1/2} \quad (5)$$

(where the characteristic age $\tau = P/2\dot{P}$) from which it follows that the lines of $E_{el} = \text{constant}$ roughly pass parallel to the lines of $\tau = \text{constant}$ on the P - $\dot{P}$ diagram for fixed values of I and R . Dependence of τ , E_{el} , B , and $\dot{E}$ on the values of I , R , and the braking index (n) and also the evolutions of pulsars with very different parameters are examined in detail by Guseinov et al. ^{6,7}.

The existing theories do not give us the possibility to estimate the value of the voltage better than the order of magnitude. But it is exactly known that charged particles can be pulled out from the hot atmosphere of pulsar because of the large value of E_{el} and they can be additionally accelerated. Further acceleration can take place in the field of the magneto-dipole wave ⁴. Which of these mechanisms is mainly responsible for the acceleration must be determined from the observational data. It may be the case that the voltage creates only basis for further acceleration of particles similar to the case of the Maxwell tail of hot particles (or particles which have already been accelerated with some other mechanism) in the shock fronts of SNRs. In the regular acceleration models ⁸⁻¹⁰, a further acceleration of particles with high energy takes place under the crossing through the shock fronts of SNRs ¹¹. Therefore, it is necessary to analyze the observational data.

3 How are the observational data related to the theoretical idea?

From the observations of the radio and X-ray radiation of neutron stars with power-law spectra (excluding the cooling radiation of neutron stars), there

is direct evidence for continuing particle acceleration. Such X-ray radiation have been observed, for example, from pulsars J1856+0113 and J1801-2451 which have PWNe and have values of $\dot{E} \sim (4-5) \times 10^{35}$ erg/s and 2.5×10^{36} erg/s, respectively. On the other hand, old millisecond pulsars J1939+2134 and J1824-2452 with $\dot{E} \sim (1-2) \times 10^{36}$ erg/s also have similar X-ray radiation as the previous pulsars², but with smaller luminosity in the 2-10 keV band. As seen in Table 1, these pulsars, beginning from J1856+0113, have values of X-ray radiation 10^{33} , 1.5×10^{33} , 5×10^{32} , and 3.6×10^{33} erg/s, respectively, in the 2-10 keV band^{2,3}. We have estimated the ratio $(\frac{\dot{P}}{P})^{1/2} \sim \tau^{-1/2}$ from the dependence of E_{el} (see expression (5)) for pulsars J1856+0113, J1801-2451, J1939+2134, and J1824-2452 and we have found the values 9×10^{-7} , 10^{-6} , 7×10^{-9} , and $3 \times 10^{-8} \text{ s}^{-1/2}$, respectively. If in all these four cases the values of radius and moment of inertia of the pulsars are approximately the same, then the values of the electric field intensity E_{el} (see expression (5)) may also have approximately the same ratios. In other words, electric fields of old millisecond pulsars are more than 2 orders of magnitude smaller compared to young pulsars with similar $L_{2-10\text{keV}}$. $\dot{E}$ values of these four pulsars are roughly on the same order of magnitude, but the initial energy of electrons is different because their E_{el} values are different. How is it possible that the $L_{2-10\text{keV}}$ values of these two types of pulsars are comparable to each other? Young pulsars have PWNe around them, i.e. an extra X-ray radiation of the electrons in the PWN comes from the source in addition to the X-ray radiation of the neutron star. These old millisecond pulsars have $\log \tau < 8.37$.

In Figure 1, dependence of the X-ray luminosity in 2-10 keV band ($L_{2-10\text{keV}}$) on the value of $\dot{E}$ for the isolated pulsars with comparably hard spectra which are located up to 6 kpc from the Sun together with 2 pulsars in the Magellanic Clouds (see Table 1) is represented. As seen in Figure 1, pulsars J1856+0113 and J1801-2451 have $\dot{E}$ values similar to isolated old millisecond pulsars J1824-2452 and J1939+2134, and their $\dot{E}$ values are more than 300 times greater than their X-ray luminosity in 2-10 keV band. Therefore, the luminosity of the power-law X-ray radiation (which includes the radiation of the relativistic particles in the Coulomb and magnetic fields) mainly depends on the value of $\dot{E}$. How can this approach be confirmed also for other pulsars based on the observational data?

Since we are interested in the acceleration of particles, we have chosen the pulsars for which $L_{2-10\text{keV}}/L_{0.1-2.4\text{keV}} > 1$, so that, we avoid the errors

connected to the interstellar absorption and the radiation related with the cooling of neutron stars. As seen from Table 1, these conditions are not satisfied for the last 4 pulsars with $\log \tau=9.58-9.86$ which are also displayed in Figure 1. But they are very close to the Sun and very old, therefore they do not create any difficulties. All the young pulsars shown in Figure 1 have not only hard spectra but also have PWNe. As seen from the equations of the best fit

$$L_{2-10keV} = 1.42 \times 10^8 \dot{E}^{0.63} \quad (6)$$

for the pulsars with $\dot{E} < 10^{35}$ erg/s and $L_{2-10keV} < 10^{31}$ erg/s and

$$L_{2-10keV} = 1.86 \times 10^{-37} \dot{E}^{1.92} \quad (7)$$

for the other pulsars with higher $L_{2-10keV}$ and/or higher $\dot{E}$, there exist clear relations between $\dot{E}$ and $L_{2-10keV}$ for which the radiation is related with the accelerated particles.

In Figure 2, the relation between $L_{2-10keV}$ and the characteristic age of the same pulsars shown in Figure 1 is represented. As seen from the figure, a single dependence for both the young and the old pulsar populations does not exist (see also Possenti et al. ³). Pulsars with similar values of $L_{2-10keV}$ and $\dot{E}$ may have very different values of characteristic age and this leads to different dependence for each population. As their ages increase, the young pulsars continue to move practically along the same direction on the L_x - τ diagram. This is seen from Table 1 where there are isolated pulsars with characteristic ages 10^5 - 10^7 yr and with luminosity in 2-10 keV band similar to the luminosity of the old millisecond pulsars. Therefore, the electric field intensity does not determine the intensity and possibly also the spectra of the relativistic particles. The X-ray luminosity strongly depends on the value of $\dot{E}$ and weakly depends on τ and E_{el} . Large values of E_{el} mainly trigger the acceleration of particles.

As seen from Figure 1 and expressions (6) and (7), as $\dot{E}$ decreases the X-ray luminosity of pulsars also decreases. So, in order to have the probability to observe X-ray radiation from a pulsar to be high for a fixed value of E_{el} , the rotation period of the pulsar must be short (see expression (4)). On the other hand, as seen from Figure 1 and Table 1, the L_x values of the young pulsars which are connected to SNRs are ~ 5 orders of magnitude larger than the L_x values of old ms pulsars and this is also seen from expressions (6) and

(7). This must be the result of small number of relativistic particles in the magnetospheres of old ms pulsars.

As seen from Figure 2, the difference in the τ values of very old ms pulsars ($\tau > 10^9$ yr) and young pulsars is about 5.5 orders of magnitude (and from expression (5) the difference in the E_{el}^2 values is also about 5.5 orders of magnitude). The differences in the $\dot{E}$ values is about 3-3.5 orders and in the P values is on average 2.5 orders of magnitude. Considering that the expression for E_{el} can only be roughly determined as mentioned above, these data show that the relation between $\dot{E}$, E_{el} and P in expression (4) is valid.

In order to tear off particles from the surface of neutron star it is necessary to have large values of E_{el} . The occurrence of this process must be easier under the existence of hot and extended atmosphere. Therefore, the values of mass and age of pulsars must also have important roles in the formation of the X-ray emission.

If the braking index n is constant along all the evolutionary tracks with different initial parameters, then the age of the pulsar is

$$t = \frac{P}{(n-1)\dot{P}} \left(1 - \left(\frac{P_0}{P}\right)^{n-1}\right) = \tau \times \left(1 - \left(\frac{P_0}{P}\right)^{n-1}\right). \quad (8)$$

For the young pulsars P_0 (the initial spin period of pulsar) can assumed to be much less than P , so that, τ can approximately be equal to the age. For recycled ms pulsars, the P value can be comparable to the P_0 value (the period which the radio pulsar is born with after completing the X-ray binary phase, see Bisnovatyi-Kogan and Komberg^{12,13} on recycled ms pulsars), so that τ can be several times larger than the age. For our purposes such a difference is not significant since it is enough to know the order of magnitude of the age which is much longer than the cooling time of recycled pulsars.

As the last 4 pulsars displayed in Table 1 with $\dot{E}$ values close to 3×10^{33} erg/s can emit X-ray, all of the isolated old ms pulsars with $\dot{E} > 3 \times 10^{33}$ erg/s must also have X-ray radiation. On the other hand, all the 5 old ms pulsars with $P \leq 6$ ms which are located up to ~ 0.5 kpc^{14,15} have been detected in X-ray^{2,3}. Since the number density of old ms pulsars is very large compared to the number density of low mass X-ray binaries, the ages of these pulsars must not be much less than τ in general. As we see, although the recycled millisecond pulsars are very old, they still emit X-ray because they have large $\dot{E}$ values.

The X-ray luminosity of pulsars together with their PWNe are represented in Table 1. Often, such X-ray luminosity values include both the X-ray radiation coming from the pulsar and the PWN together, because it is difficult to separate the emission of the pulsar from the emission of the PWN. Therefore, the radiation of isolated ms pulsars must be smaller compared to isolated young pulsars for equal values of $\dot{E}$. This must be true also because of young pulsars having large values of electric field intensity and possibly having smaller masses in some cases. Therefore, it is strange that ms pulsar J1824-2452 has larger luminosity compared to young pulsar J1801-2451 (see Figure 1). On the other hand, their radiation is very hard. Actually, the uncertainties in the luminosity values of these ms pulsars may be large.

In Figure 3, P- $\dot{P}$ diagram of all the pulsars in Table 1, which have connections with SNRs and/or from which X-ray radiation has been observed, is displayed. As seen from this figure and Table 1, pulsars Geminga and J0538+2817 which is connected to SNR G180.0-1.7¹⁶ have about one order of magnitude larger values of $\dot{E}$ compared to old ms pulsars, but their luminosity in 2-10 keV band is similar to the luminosity of these ms pulsars. The X-ray spectra of the mentioned young pulsars are steeper compared to the spectra of old ms pulsars as seen from the comparison of the luminosity in 0.1-2.4 keV and 2-10 keV bands. The luminosity of J0826+2637 and J0953+0755 in 2-10 keV band are smaller than the luminosity of old ms pulsars, but their $\dot{E}$ values are also smaller (see Fig.3 and Fig.1). How can this observational fact be explained?

Young pulsars have values of electric field intensity about 50 times larger than the values of old ms pulsars, they are hotter and they may also have smaller masses. These facts create conditions for tearing off charged particles more easily from neutron stars. On the other hand, the magnetic field values of young pulsars are more than 3 orders of magnitude larger (the high magnetic field does not let the particles to escape from the magnetosphere without considerable radiative energy loss). These and the possible existence of large number of charged particles in the magnetospheres and the surroundings of young pulsars must create the best conditions for magneto-braking and Coulomb radiation. Among young pulsars, the small values of X-ray radiation of J0538+2817 and Geminga in 2-10 keV band seem to contradict this natural discussion.

4 Pulsar wind nebulae and pulsars in super-nova remnants

Let us now examine the presence or absence of PWN around pulsars and the X-ray radiation. All the pulsars with $\tau < 10^7$ yr represented in Table 1 (except pulsar J0537-6910 in the Magellanic Cloud) are shown on the P- $\dot{P}$ diagram in Figure 4 denoting also the morphological types of the SNRs which some of the pulsars are connected to. If the type of the SNR is C (composite) or F (filled-center), then it means that there is PWN created by the neutron star. In S (shell) type SNRs there is no observed PWN possibly because it is very faint. Other pulsars in Table 1 which are not connected to SNRs and which have $\tau < 10^7$ yr are also displayed in Figure 4. All the pulsars with PWN around them have L_x (2-10 keV) $> 10^{33}$ erg/s (see Table 1). The cooling radiation of these pulsars do not have a considerable role.

Pulsars J1803-2137, J1016-5857, J2337+6151, and J0659+1414, which are connected to S-type SNRs (in the radio band), have been observed to radiate in X-ray band. Pulsars J2337+6151 and J0659+1414 have suitable distance values and positions to observe possible PWNe around them, but $\dot{E}$ and L_x (2-10 keV) values of these pulsars are so small that it is not so likely that PWNe with observable brightness are present around them.

From the analysis of these data, we see that lifetime of the X-ray radiation produced by the relativistic particles in the magnetospheres of neutron stars is considerably longer than lifetime of PWNe (see Table 1). But there may be one exception, namely pulsar J0538+2817, which is connected to SNR G180.0-1.7 (S147). This SNR is S-type in radio band ¹⁷. Romani & Ng ¹⁶ claim evidence for a faint nebula around the pulsar, but in a more recent work ¹⁸ no evidence has been found of a PWN and the pulsar has been observed to radiate only thermally. Also, the position and ages of the pulsar and the SNR show that a physical connection between these two sources is dubious and there are no data directly showing evidence for the connection.

From Figure 4 and Table 1 we see that PWN may exist around pulsars with $\dot{E} > 10^{35}$ erg/s and $\tau < 5 \times 10^4$ yr. Pulsar J1646-4346 in C-type SNR G341.2+0.9 satisfies these criteria, but X-ray radiation has not been observed from it because it is located at a distance of 6.9 kpc ¹⁹. X-ray radiation has not been observed also from 9 far away pulsars with $d \geq 3$ kpc which are located close to the Galactic plane in the Galactic central directions (see Table

1 and Figure 4) and which satisfy the conditions of large value of $\dot{E}$ and small value of τ . As seen from the position of pulsar J1617-5055 in Figure 4, it must have a connection with SNR and must have PWN. This follows also from the large values of X-ray luminosity in both X-ray bands. Lack of observed SNR and PWN must be related with the large distance and the position in the Galaxy. Ages of the relatively nearby pulsars which are displayed in Figure 4 may be smaller than the values of τ . Three of these pulsars are connected to S-type SNRs and all of them have small X-ray luminosity.

5 Conclusions

From the analysis of all the available data on isolated X-ray pulsars, their wind nebulae and the SNRs which are connected to some of these sources and also from theoretical considerations, we have found the conclusions below:

1. The electric field intensity (voltage) of neutron stars is not the main physical quantity for the spectra, energy, and intensity of the relativistic particles which produce the X-ray radiation of isolated pulsars and their wind nebulae. The voltage mainly triggers the acceleration of particles. In order to tear off charged particles from the surface of neutron star E_{el} must be large. This process occurs more easily in hot and extended atmospheres. Therefore, the values of mass and age of pulsars also have important roles in the formation of the X-ray radiation.
2. The acceleration of relativistic particles mainly takes place in the field of the magneto-dipole radiation wave. Because of this, the X-ray radiation of pulsars and their wind nebulae strongly depend on the value of the rate of rotational energy loss which is reflected by the spectra of the magneto-dipole radiation.
3. The high magnetic field and particularly the number density of the charged particles create conditions under which the energy loss of the particles in 0.1-2.4 keV band may surpass the energy received from the magneto-dipole radiation wave for further acceleration. This must be responsible for the steeper X-ray spectra of young pulsars as compared to old ms pulsars.
4. PWN exists only around pulsars with $\dot{E} > 10^{35}$ erg/s and $\tau < 5 \times 10^4$ yr. Also, the X-ray luminosity of pulsars in 2-10 keV band drops down to $\sim 10^{33}$ erg/s at about the same time when the PWN becomes unobservable in both radio and X-ray bands. The lifetime of S-type SNRs exceeds the lifetimes

of F-type and C-type SNRs. So, it may be possible that C-type SNRs can show themselves as S-type after some time in their evolution. PWN must be observed around pulsar J1617-5055.

Acknowledgments S.O.T is supported by INTAS. A.A. and A.M.A. thank Bogazici University Foundation (BUVAK).

References

1. W. Becker and J. Trumper, *Astron. Astrophys.* **326**, 682 (1997).
2. W. Becker and B. Aschenbach, Proceedings of the 270. WE-Heraeus Seminar on: "Neutron Stars, Pulsars and Supernova Remnants", eds. W. Becker, H. Lesch and J. Trumper, MPE Report 278, p.64 (2002).
3. A. Possenti, R. Cerutti, M. Colpi and S. Mereghetti, *Astron. Astrophys.* **387**, 993 (2002).
4. V. M. Lipunov, *Astrophysics of Neutron Stars*, Springer-Verlag (1992).
5. A. G. Lyne and F. Graham-Smith, *Pulsar Astronomy*, Cambridge University Press (2006).
6. O. H. Guseinov, A. Ankay, S. O. Tagieva, *Int. J. Mod. Phys. D* **14**, 1465 (2005).
7. O. H. Guseinov, A. Ankay, S. O. Tagieva, *Astrophys. Space Sci.* **298**, 553 (2005).
8. A. R. Bell, *Mon. Not. Roy. Astron. Soc.* **182**, 147 (1978).
9. A. R. Bell, *Mon. Not. Roy. Astron. Soc.* **182**, 443 (1978).
10. G. F. Krymsky, *Dokl. Akad. Nauk SSSR* **234**, 1306 (1977).
11. A. O. Allakhverdiev, A. I. Asvarov, O. H. Guseinov, F. K. Kasumov, *Astrophys. Space Sci.* **123**, 237 (1986).
12. G. S. Bisnovaty-Kogan and B. V. Komberg, *Sov. Astron.* **18**, 217 (1974).
13. G. S. Bisnovaty-Kogan and B. V. Komberg, *Sov. Astron. Lett.* **2**, 130 (1976).
14. ATNF – Australia Telescope National Facility Pulsar Catalogue (2007), <http://www.atnf.csiro.au/research/pulsar/psrcat/>.
15. O. H. Guseinov, S. K. Yerli, S. Ozkan, A. Sezer and S. O. Tagieva, *Astron. Astrophys. Transac.* **23**, 357 (2004).
16. R. W. Romani and C. -Y. Ng, 2003, *Astrophys. J.*, **585**, L41 (2003).
17. D. A. Green, 'A Catalogue of Galactic Supernova Remnants (2006 April version)', (<http://www.mrao.cam.ac.uk/surveys/snrs/>).
18. K. E. McGowan, J. A. Kennea, S. Zane, et al., *Astrophys. J.* **591**, 380 (2003).
19. O. H. Guseinov, A. Ankay, A. Sezer, S. O. Tagieva, *Astron. Astrophys. Transac.* **22**, 273 (2003).
20. D. J. Helfand, B. F. Collins and E. V. Gotthelf, *Astrophys. J.* **582**, 783

- (2003).
21. D. R. Lorimer, A. G. Lyne and F. Camilo, *Astron. Astrophys.* **331**, 1002 (1998).
 22. A. O. Allakhverdiev, M. A. Alpar, F. Gok and O. H. Guseinov, *Turkish J. Phys.* **21**, 688 (1997).
 23. F. D. Seward, F. R. Jr. Harnden and D. J. Helfand, *Astrophys. J.* **287**, L19 (1984).
 24. R. Kothes, B. Uyaniker and S. Pineault, *Astrophys. J.* **560**, 236 (2001).
 25. R. N. Manchester, J. F. Bell, F. Camilo, et al., Neutron Stars in Supernova Remnants, ASP Conference Series, edited by Patrick O. Slane and Bryan M. Gaensler, **271**, p.31 (2002).
 26. S. E. Thorsett, W. F. Brisken and W. M. Goss, *Astrophys. J.* **573**, L111 (2002).
 27. D. A. Frail, W. M. Goss and J. B. Z. Whiteoak, *Astrophys. J.* **437**, 781 (1994).
 28. F. Camilo, J. F. Bell, R. N. Manchester, et al., *Astrophys. J.* **557**, L51 (2001).
 29. V. M. Kaspi, M. E. Roberts, G. Vasisht, E. V. Gotthelf, M. Pivovarovoff and N. Kawai, *Astrophys. J.* **560**, 371 (2001).
 30. E. Furst, W. Reich and J. H. Seiradakis, *Astron. Astrophys.* **276**, 470 (1993).
 31. Y. Gupta, D. Mitra, D. A. Green and A. Acharyya, astro-ph/0508257.
 32. M. Kramer, A. G. Lyne, G. Hobbs, O. Lohmer, P. Carr, C. Jordan and A. Wolszczan, *Astrophys. J.* **593**, L31 (2003).

Table 1 - All 35 radio and X-ray pulsars with detected X-ray radiation in 0.1-2.4 keV and/or in 2-10 keV bands and with $\tau < 10^6$ yr. All 15 pulsars which have $\text{Log } \dot{E} > 35.6$ and $\tau < 10^5$ yr. All pulsars which are connected to SNRs (PWNe). Two of these pulsars are in Magellanic Clouds and the others are Galactic pulsars with $d \leq 7$ kpc. In the first column 'G' denotes strong glitch. In column 12 $\beta = 2\Delta\theta/\theta$ (θ : SNR diameter, $\Delta\theta$: angular distance of pulsar from the geometric centre of SNR).

JName	Type	d kpc	P s	$\dot{P}$ s/s	Log τ	Log B	Log $\dot{E}$	SNR + l,b	Type	β	Log L_x 0.1-2.4 keV	Log L_x 2-10 keV	n
J1852+0040	X	7	0.1049					G33.6+0.1	S			33.6	
J1846-0258	X	6.7	0.325	7.1E-12	2.859	13.70	36.9	G29.7-0.3	C	$\sim 0^{20}$	~ 35.18	35.32	2.65 $\pm$ 0.01
J0534+2200	ROXG	2.0	0.034	4.21E-13	3.10	12.58	38.64	Crab	F	$\sim 0.19^{21}$	35.98	36.65	2.51
J1513-5908	RXG	4.2	0.151	1.54E-12	3.191	13.19	37.25	G320.4+1.2	C	0.24^{21}	34.25	35.32	2.839
J1119-6127	RX	7.0	0.408	4.02E-12	3.205	13.61	36.37	G292.2-0.5	C	$\sim 0^{22}$	≤ 32.5	33.42	2.91
J0540-6919	ROX	50	0.051	4.79E-13	3.22	12.70	38.17	N158A	C	$\sim 0^{23}$	36.21	36.93	2.140 $\pm$ 0.009
J1124-5916	RX	6.0	0.135	7.45E-13	3.458	13.01	37.07	G292.0+1.8	C		~ 32.7	34.67	
J1930+1852	RX	7	0.137	7.51E-13	3.462	13.0	37.23	G54.1+0.3	F		~ 34.2		
J0537-6910	X	50	0.016	5.1E-14	3.7	12.0	38.7	N157B	F		~ 36.0	36.11	$> 6.9?$
J1833-1034 ³¹	R	5.5	0.061	2.0E-13	3.7	12.56	37.5	G21.5-0.9	C				
J0205+6449	RX	3.2	0.066	1.9E-13	3.74	12.55	37.42	G130.7+3.1	F		32.38	34.26	
G J1617-5055	RX	6.2	0.069	1.371E-13	3.903	12.49	37.21	332.5,-0.27			34.30	34.59	
J2229+6114	RXG	5.5	0.052	7.8E-14	4.023	12.31	37.34	G106.6+2.9	C	0^{24}	33.58	33.69	
G J0835-4510	ROXG	0.40	0.089	1.25E-13	4.05	12.53	36.84	Vela	C	0.29,0.3 ^{21,22}	32.46	33.18	1.4
J1420-6048	RXG	6.1	0.068	8.32E-14	4.113	12.38	37.01	G313.4+0.2?	F	0.2^{25}	34.26	34.30	
G J1801-2451	RX	4.5	0.125	1.28E-13	4.19	12.61	36.41	G5.27-0.9	F	$\sim 0^{26}$	~ 33.0	33.18	
G J1803-2137	RX	3.5	0.134	1.34E-13	4.197	12.63	36.35	G8.7-0.1	S?	0.7^{27}	33.06	32.39	
J1702-4310	R	4.8	0.2405	2.24E-13	4.23	12.88	35.80	343.4,-0.85					
G J1709-4428	RXG	1.8	0.102	9.30E-14	4.241	12.49	36.53	G343.1-2.3?			33.15	32.58	
J1856+0113	RX	2.8	0.267	2.08E-13	4.31	12.88	35.63	G34.7-0.4	C	$0.51^{21}, 0.6^{22}$	≤ 33.0	33.03	
J1048-5832	RXG	2.8	0.124	9.63E-14	4.31	12.54	36.30	287.4,+0.58			≤ 32.11	32.35	
J1016-5857	RX	6	0.107	8.11E-14	4.321	12.48	36.41	G284.3-1.8	S	$1^{25}, 1.3^{28}$	~ 32.6		
G J1826-1334	RX	3.4	0.101	7.55E-14	4.33	12.45	36.45	G18.0-0.7	F		~ 32.7	34.34	
J1811-1925	X	5	0.065	4.22E-14	4.4	12.23	36.78	G11.2-0.3	F	$\sim 0^{29}$	32.8	34.54	
J1747-2958	RX	2.0	0.099	6.14E-14	4.41	12.41	36.41	G359.2-0.8?	F		~ 34.0		
G J1730-3350	R	4.24	0.1394	8.51E-14	4.41	12.56	36.09	354.1,+0.09					
J1646-4346	R	6.9	0.232	1.13E-13	4.51	12.71	35.55	G341.2+0.9	C	0.7^{27}			
J1837-0604	R	6.2	0.0963	4.52E-14	4.53	12.34	36.30	26.0,+0.27					
J1015-5719	R	4.87	0.1399	5.74E-14	4.59	12.47	35.92	283.1,-0.58					
J1718-3718	RX	4.5	3.3	1.5E-12	4.6	13.9	33.2	349.85,+0.22			33.8	33? ³²	
J2337+6151	RX	2.8	0.495	1.92E-13	4.611	12.99	34.79	G114.3+0.3	S	$0.08^{21,30}$	31.90	31.47	
J1637-4642	R	5.5	0.1540	5.92E-14	4.62	12.50	35.81	337.8,+0.31					
J0940-5428	R	3.8	0.0875	3.29E-14	4.63	12.25	36.29	277.5,-1.29					
J0631+1036	RX	6.6	0.288	1.05E-13	4.639	12.74	35.24	201.2,+0.45				31.90	
J1809-1917	R	3.3	0.0827	2.55E-14	4.71	12.18	36.25	11.1,+0.08					
J1105-6107	RX	7.0	0.063	1.58E-14	4.801	12.01	36.39	290.5,-0.85			~ 33.2	33.55	
J1718-3825	R	3.3	0.0747	1.32E-14	4.95	12.02	36.10	348.9,-0.43					
J1531-5610	R	2.6	0.0842	1.37E-14	4.99	12.05	35.96	323.9,+0.03					
J1952+3252	RXG	2.0	0.040	5.84E-15	5.030	11.69	36.57	G69.0+2.7	C	0.14,0.15 ^{21,22}	33.64	33.07	
J0659+1414	ROX	0.3	0.385	5.50E-14	5.044	12.67	34.58	G201.1+8.7			32.18	30.32	
J0908-4913	R	4.5	0.1068	1.51E-14	5.05	12.12	35.69	270.3,-1.02					
J0855-4644	R	6.4	0.0647	7.26E-15	5.15	11.85	36.03	267.0,-1.0					
G J1833-0827	R	5.67	0.0853	9.17E-15	5.17	11.97	35.77	23.4,+0.06					
J1509-5850	R	2.5	0.0889	9.17E-15	5.19	11.97	35.71	320.0,-0.62					
J1913+1011	R	4.1	0.0359	3.37E-15	5.23	11.56	36.46	44.5,-0.17					

Table 1 - continued

	JName	Type	d kpc	P s	$\dot{P}$ s/s	Log τ	Log B	Log $\dot{E}$	SNR + l,b	Type	β	Log L_x 0.1-2.4 keV	Log L_x 2-10 keV	n
	J0117+5914	RX	2.4	0.101	5.85E-15	5.44	11.89	35.34	126.3,-3.46				30.44	
Be	J1302-6350	RX	1.3	0.048	2.28E-15	5.52	11.52	35.92	304.2,-1.00			32.95	32.57	
	Geminga	X	0.15	0.237	1.14E-14	5.53	12.22	34.53	195.13,+4.27			31.10	29.33	
	J1057-5226	RX	1.0	0.197	5.83E-15	5.73	12.04	34.48	286.0,+6.65			33.13	30.08	
G	J0358+5413	RX	2	0.156	4.4E-15	5.75	11.92	34.66	148.2,+0.81			31.96	31.75	
	J0538+2817	RX	1.5	0.143	3.67E-15	5.79	11.87	34.69	G180.0-1.7?	S	0.41 ³²	32.74	29.31	
	J1932+1059	RX	0.2	0.227	1.16E-15	6.49	11.71	33.59	47.38,-3.88			30.07	29.41	
	J0826+2637	RX	0.4	0.531	1.71E-15	6.69	11.98	32.65	196.96,+31.74			31.85	29.01	
	J0953+0755	RX	0.13	0.253	2.29E-16	7.24	11.39	32.75	228.91,+43.7			29.35	28.62	
	J1824-2452	RX	5.1	0.003	1.62E-18	7.48	9.35	36.35	7.8,-5.58			32.2	33.56	
	J1939+2134	RX	3.6	0.0015	1.06E-19	8.37	8.61	36.04	57.51,-0.29			<32.1	32.54	
	J2124-3358	RX	0.25	0.005	2.0E-20	9.58	8.51	33.83	10.93,-45.44			30.35	29.58	
	J1024-0719	RX	0.35	0.005	1.9E-20	9.64	8.50	33.73	251.7,+40.52			29.48	29.09	
	J0030+0451	RX	0.23	0.005	1.0E-20	9.89	8.35	33.53	113.14,-57.61			30.56	29.88	
	J1744-1134	RX	0.36	0.004	8.9E-21	9.86	8.29	33.72	14.79,+9.18			29.58	29.25	

Figure Captions

Figure 1: Dependence of the 2-10 keV luminosity on the rate of rotational energy loss for pulsars located up to 7 kpc from the Sun including also 2 pulsars in Magellanic Clouds which are denoted by '+' sign. Old millisecond pulsars are shown by 'star' sign and young pulsars by 'X' sign.

Figure 2: Dependence of the 2-10 keV luminosity on the characteristic age for pulsars located up to 7 kpc from the Sun including also 2 pulsars in Magellanic Clouds. The pulsars are displayed by the same symbols as in Figure 1.

Figure 3: Period versus period derivative diagram for the pulsars which have been detected in X-ray and/or which are connected to SNRs (see Table 1). The sign '+' represents the pulsars which have physical connections with SNRs and the 'X' sign denotes the pulsars which are not connected to SNRs. B8-B14, E29-E41 and T3-T10 represent the constant lines of the effective surface magnetic field ($B=10^8$ - 10^{14} Gauss), the rotational kinetic energy loss ($\dot{E}=10^{29}$ - 10^{41} erg/s) and the characteristic age ($\tau=10^3$ - 10^{10} yr).

Figure 4: Period versus period derivative diagram for all 48 pulsars represented in Table 1 with $\tau < 10^6$ yr and distance ≤ 7 kpc including the 2 pulsars in Magellanic Clouds. '+' signs represent the 11 pulsars connected to C-type SNRs, 'X' signs show the 9 pulsars connected to F-type SNRs, and 'star' signs denote the 4 pulsars connected to S-type SNRs. 'Light square' signs represent the 14 pulsars which are not connected to SNRs and not observed in X-ray. The 10 pulsars which are denoted with 'dark squares' have been observed in X-ray but they are not connected to SNRs. B12-B14, E32-E38 and T3-T6 represent the constant lines of the effective surface magnetic field ($B=10^{12}$ - 10^{14} Gauss), the rotational kinetic energy loss ($\dot{E}=10^{32}$ - 10^{38} erg/s) and the characteristic age ($\tau=10^3$ - 10^6 yr).







